Pend Orielle County Diabetes Statistics, 2003



 Prevalence
 Pend Orielle County
 Washington State

 (adults 18+ years)
 Percent (95% CI)¹ Number of adults
 Percent (95% CI)¹ Number of adults

 13.0 (6.8, 23.5)
 1,418
 6.6 (6.2, 7.1)
 303,283

Diabetes Prevention & Control Program P.O. Box 47836 Olympia, WA 98504-7836 (360) 236-3617

Risk Factors	Pend Orielle County	Washington State
(adults 18+ years without diabetes)	Percent (95%CI) ¹	Percent (95%CI) ¹
Overweight or obese ²	65.8 (58.0, 72.8)	56.8 (55.8, 57.8)
Insufficient physical activity ³	34.3 (26.8, 42.8)	35.8 (34.9, 36.8)
Smoke cigarettes	28.1 (19.6, 38.6)	19.8 (19.0, 20.6)
History of high blood pressure ⁴	28.4 (22.1, 35.7)	20.8 (20.1, 21.5)
History of high cholesterol ⁴	41.9 (34.3, 50.0)	31.1 (30.2, 32.1)

	Number of	Age-adjusted rate per 10,000 total		Crude rate per 10,000 people with	
Hospitalizations ⁵	hospitalizations	population (95% CI) ¹		diabetes	
(with complications of diabetes)	Pend Orielle County	Pend Orielle County	Washington State	Pend Orielle County	Washington State
Diabetes as first listed diagnosis	14	10.6 (5.7, 19.4)	9.6 (9.4, 9.9)	98.2	191.3
Diabetes as any listed diagnosis	174	125.1 (106.7, 146.7)	118.4 (117.5, 119.3)	1,220.3	2,293.6
→ with coronary heart disease	46	32.9 (23.8, 45.4)	36.3 (35.8, 36.8)	322.6	693.1
→ with stroke	11	7.7 (3.8, 15.5)	10.1 (9.9, 10.4)	77.1	190.9
with diabetic ketoacidosis	*	*	3.9 (3.8, 4.1)	*	78.9
with lower extremity amputation	5	3.6 (1.1, 10.3)	1.9 (1.8, 2.0)	35.1	160.7

Deaths ⁵	Number of deaths	Age-adjusted rate per 100,000 total population (95% CI) 1		Crude rate per 100,000 people with diabetes	
	Pend Orielle County	Pend Orielle County	Washington State	Pend Orielle County	Washington State
Diabetes as underlying cause	7	45.7 (18.0, 113.1)	26.0 (24.7, 27.4)	490.9	494.4
Diabetes as any cause	17	109.4 (63.1, 191.2)	77.1 (74.9, 79.4)	1,192.2	1,460.9

¹Data presented as: percentage or rate (95% confidence interval).

Please see http://www.doh.wa.gov/cfh/diabetes/publications/Face%20Page.pdf for a description of data sources and methods used to develop these county profiles (includes estimates of total diabetes population for each county that are used to compute the hospitalization and death rates per people with diabetes).

²Overweight=body mass index (BMI) 25.0-29.9 kg/m², obese=BMI 30.0+ kg/m².

³Insufficient physical activity=reported level of physical activity during leisure time or at work does not meet Centers for Disease Control and Prevention (CDC) recommendations of moderate-intensity activity for 30+ minutes on 5+ days/week or vigorous-intensity activity for 20+ minutes on 3+ days/week

⁴Told by doctor, nurse, or other health professional had high blood pressure and/or high cholesterol.

⁵Rates are not computed for counts <5. Rates based on fewer than 20 deaths or hospitalizations are likely to be unstable or imprecise.